Urban Aerosol-Induced Changes of Precipitation

Daniel Rosenfeld, Amir Givati,
Alexander Khain, Guy Kelman
Earth Sciences, The Hebrew University of Jerusalem, Israel

and


William L. Woodley Woodley Weather Consultants




Effect of Added CCN

(Cloud Condensation Nuclei)

- Adding CCN makes clouds with more, smaller droplets.
- These clouds are whiter, reflect more sunlight
- Too many tiny CCN suppress coalescence and precipitation




Industrial pollution tracks over Manitoba, Canada





Ship tracks off the Washington coast

Tropical Rainfall Measuring Mission (TRMM) satellite observations show how rain is inhibited in clouds forming in dirty air, i.e., air containing many small CCN, contributed by aerosol particles such as smoke, air pollution and desert dust.




Rosenfeld D., 1999: TRMM Observed First Direct Evidence of Smoke from Forest Fires Inhibiting Rainfall. *Geophysical Research Letters.* **26**, (20), 3105-3108.







Suppression of Rain and Snow by Urban and Industrial Air Pollution (Rosenfeld, 2000, Science)




VIRS painting yellow pollution tracks in the clouds over South Australia, due to reduced droplets size. PR shows precipitation as white patches only outside the pollution tracks, although clouds have same depth.




VIRS retrieved effective radius does not exceed the 14 µm precipitation threshold in polluted clouds within area 2 in the Australia image.



PR shows bright band in clean clouds. Therefore, pollution suppressed rain and snow in polluted clouds.










Table 1. Chemical and Physical Properties of Cloud Droplets and Snow During Two Precipitation Events

February	15	19	
Major Habit	Planar Dendrite	Planar Dendrite	
Rime Category	Unrimed (0.5)	Moderate (2.0)	
Rime Mass Frac.	5%	51%	
SPL Precip. Rate	0.02 mm hr^{-1}	0.38 mm hr^{-1}	
ISS Precip.Rate	0 to 0.1 mm hr^{-1}	1.1 mm hr^{-1}	
SPL Temperature	−13°C	$-4^{\circ}C$	
Snow δ ¹⁸ O	-22.1	-16.5	
Cloud δ ¹⁸ O	-21.1	-16.2	
δ ¹⁸ O Snow Mass	−14°C	-4.8° C	
Temp. Of Origin			
Cloud Top Temp	−19°C	−22°C	
Snow CAE SO ₄	$0.011 \ \mu g \ m^{-3}$	$0.072~{\rm \mu g~m^{-3}}$	
Cloud CAE SO ₄	1.1 $\mu g m^{-3}$	$0.12 \ \mu g \ m^{-3}$	
Droplet Mean Dia.	8.3 mm	13.6 mm	
Droplet Conc.	310 cm^{-3}	74 cm^{-3}	
Cloud SCLW	0.13 g m^{-3}	0.14 g m^{-3}	

Figure 4. Cloud droplet and snow size spectra for the sampling periods on Feb. 15 and 19, 2001.

on Feb. 15 and 19 were 0.02 mm hr⁻¹ and 0.38 mm hr⁻¹, respectively.

The difference between the cloud clear air equivalent anthropogenic aerosol sulfate concentrations on the two days is nearly an order of magnitude, but in absolute terms it is only 1 mg m⁻³. Astonishingly, this small amount of aerosol can reduce the snowfall rate up to 50%.

Evidence is presented to demonstrate the possible magnitude of the secondary indirect aerosol effect on precipitation rates from cold mixed-phase clouds in mountainous regions where a seeder-feeder cloud couplet is present. Changes as small as 1 μ g m⁻³ in CCN aerosol concentration can cause significant changes in cloud properties and precipitation efficiencies. (Quoted from Borys et al., GRL 2003).


The study purposes:

- Quantify the effects of air pollution on precipitation downwind of major urban areas
- Determine the conditions in which the pollution effects are mostly effective

The study principles:

- The effect will be shown best where maritime air is polluted over coastal urban areas, and the polluted air rises over mountains downwind and forms new polluted clouds.
- The effect will be manifested as a reduction of the orographic enhancement factor with respect to the upwind coastal rainfall.
- The effect is most detectable in highly correlated mountain and coastal rain stations.

No effect is expected in nearby rural mountain-coast stations.







Legend

1E. Ukiah

2E. Lake Spaulding

3E. Bowman

4E. Boca

5A. San Francisco

6A. Sacramento

7A. Pacific House

8A. Cluster of snow packs in the divide line downwind to Sacramento 9A. Woodfords

10B. Fresno

11B. Grant Grove

12B. Cluster of snow packs in the divide line downwind

to Fresno

13B. Glacier

14B. Bishop Lake

15F. Lompoc

16F. Mt. Figuroa

17F. Santa Barbara

18F. Mt. pine

19C. Los Angeles area

20C. Cluster of stations downwind to Los Angeles


21C. Lake Arrowhead

22C. Big Bear Lake


23C. Morongo


24. San Diego


25. Cuyamaca
















Why use the ratio between two stations?


Both the plain (Israel central coast) and the Hill stations (Judea Hills) show an increase in the yearly rain amounts





But the ratio (the orographic component of the precipitation) between the hill to the plain stations is decreasing!

The Ratio between cluster of stations in Judea hills to cluster of station in the Israel plain Ending / starting ratio = 1.19 / 1.38 = 0.84, Pvalue = 0.0006




Same thing happens in southern California: The plains and the Hill stations show an increase in the yearly rain amounts

Plain stations:

Los Angeles 355, Los Angeles CC, Pomona, Los Angeles AP, Beverly Hills 22, Chino, San Bernardino


Yearly avarage rain - Cluster of stations in L.A plain area




Mountain stations:

Lake Arrowhead, Sierra PH, RaywoodFlats, Crystal

Yearly avarage rain - Cluster of Mountain stations in L.A area



But the ratio between the mountain and the plains stations is decreasing




The study areas:


California coastal range and the Sierra Nevada


Israel central mountain range


<u>Legend</u>


- 1E. Ukiah
- 2E. Lake Spaulding
- 3E. Bowman
- 4E. Boca
- 5A. San Francisco
- 6A. Sacramento 7A. Pacific House
- 8A. Cluster of snow packs in the divide line downwind
- to Sacramento 9A.Woodfords
- 10B. Fresno
- 11B. Grant Grove
- 12B. Cluster of snow packs
- in the divide line downwind
- to Fresno
- 13B. Glacier
- 14B. Bishop Lake
- 15F. Lompoc
- 16F. Mt. Figuroa
- 17F. Santa Barbara
- iri. Saika Daiba
- 18F. Mt. pine
- 19C. Los Angeles area
- 20C. Cluster of stations
- downwind to Los Angeles
- 21C. Lake Arrowhead
- 22C. Big Bear Lake
- 23C. Morongo 24. San Diego
- 25. Cuvamaca



























Legend 1E. Ukiah 2E. Lake Spaulding 3E. Bowman 4E. Boca 5A. San Francisco 6A. Sacramento 7A. Pacific House 8A. Cluster of snow packs in the divide line downwing		HOOPAH (VOLLEYUR Shasta Lak Redding	Mt. L	LAKE DESERT A S II N A D A			
to Sac 9A.Wo 10B. F 11B. O 12B. O in the to Fre 13B. O 14B. E 15F. L	cramento codfords Fresno Grant Grove Cluster of snow packs e divide line downwind	Santa Rosa⊚ Vallaji San Francisco Haywa San J	Californ	ot of the or	A Telesque	lonic con precipitat by the Natio Atmosph	nal
17F. S 18F. N 19C. I		Elev.(m)	Period	Mean μeq I ⁻¹	2001- 2002	Trend μeq I-y-1	Precip. Ro change %
20C. (down 21C. I 22C. I	Sequoia/ Fresno	1902	1981- 2002	41.3	35.4	+0.31	-24 (1945- 2000)
23C. I 24. Sa 25. C		1408	1982- 2002	30.7	36.5	-0.06	-22 (1945- 2000)
	Lassen/ Ukiah	1765	2001- 2002		17.7		0 (1945- 2000)














Analysis of the orographic factor trends according to the synoptic conditions:

- 3°C : Expected Clouds Temperatures at 700 mb on rainy days - Los Angeles and San Diego areas



The annual ratios of precipitation (Ro) between Cuyamaca and San Diego for clouds occurring when T>-3°C at 700 hpa (mainly frontal and warm air mass) and when $T \le -3$ °C (mainly cyclonic post frontal clouds).





The radiosonde model

In order to separate and identify the human potential causes from natural processes, such as changes in the atmospheric circulation, a model that predicts the natural rain in the mountain was calculated:

RMM = (RCM*X1) + (WCOMP*W*X2) + Const

RMM is the predicted precipitation In the mountains

RCM is the gauged precipitation at the coast

WCOMP is the wind speed component toward the mountain (850 mb)

W is the mixing ratio (850 mb)


Const is the multiple regression constant

The radiosonde model results - California:


The ratio between measured daily precipitation to the model-predicted daily precipitations (R=0.80 and 0.75) in polluted areas (San Diego and Los Angeles areas)

The ratio between measured daily precipitation to the model predicted daily precipitation (R=0.86) in a "clean" area (Monterey county)






--⊽-- Measured / Model





The radiosonde model results - Israel:


The ratio between measured daily precipitation in the **Judea** hills to the model predicted daily precipitation (R = 0.87) in a polluted area

































Topographic cross section showing the effects of urban air pollution on precipitation as the clouds move from west to east across the Sierra Nevada





Conclusions:

- All time series from metropolitan areas in California show a decrease in the ratio between the mountain and coastal stations along the years (mostly between 1950 to 1980). Similar trends were found also in Israel
- In the control areas, no change at all in the orographic enhancement factor was detected, both in California and in Israel
- The decrease was found only in cold air masses with clouds ingesting the air from the polluted boundary layer.
- Based on the sounding analyses, no evidence was found that can explain natural trends in the orographic enhancement factor in California and Israel
- The likely explanation for the 15%-25% decreasing orographic rainfall is the air pollution

Research Recommendations

Satellite Microphysical Survey

Numerical Modeling with Explicit Microphysics and Varying Aerosol Inputs

Measurements of Aerosol Burden and Transport in California

Analyses to Separate Out Effects of Glaciogenic Seeding from Effects of Pollution


Additional Measurements of the Effects of Anthropogenic Aerosols on Snow Growth and Snowfall Rate

Develop Relationships Relating Precipitation to Runoff


Dr. Woodley's formal presentation ended at this point. The following slides have been left in this presentation for those who are interested in more information on this subject.



Continental: Polluted, Suppressed rain, Strong updraft



Maritime: Clean, Fast rain, Suppressed updraft



- Cloud drop
- Rain drop
- Ice crystal
- O Ice precipitation



Bangkok Sumatra Kalimantan 🎝 10 15

The effect of aerosols on precipitation in clouds was calculated from the data of the image above. The warm colors represent efficient precipitation processes, while the cold colors represent suppressed precipitation, due to the pollution. The scale is the maximal cloud top temperature [°C] required for onset of precipitation.

(1) Maritime and Rural aerosols

Clouds from *clean maritime* air develop precipitation efficiently. After interacting with *rural aerosols*, the clouds are less efficient in developing precipitation.


(2) Urban air pollution

The **blue color** indicates detrimental effect of **urban air pollution** on the precipitation in the clouds.

(3) Smoke from forest fires

Another case of detrimental effect of the interaction of Clouds with biomass burning smoke on the precipitation in the clouds can be seen in the **blue color** over Sumatra and Kalimantan.

The TOMS aerosol index can be seen below:




So, does air pollution suppress or enhance overall rainfall amount from convective clouds?

Observations and model simulations show that always clouds with more small CCN will rain less for a given maximum vertical development.

Simulations show that in warm base clouds elevating the onset of precipitation can lead to longer time of cloud growth before downdrafts take over, and hence this dynamic feedback causes greater vigor and secondary formation of clouds, leading to more overall precipitation.


Simulation of extremely continental high base (11°C) clouds (West Texas, August 1999)

The "polluted" Cb can have nearly adiabatic water until -38°C Therefore, it has very low precipitation efficiency.




letters to nature


Deep convective clouds with sustained supercooled liquid water down to -37.5 °C




Daniel Rosenfeld* & William L. Woodley†


Nature, 25 May 2000, vol 405, p. 440-442.






West Texas: rain accumulation









Prestorm Alabama:


Time evolution of accumulated rain



PRESTORM: MAX. and MIN. VERTICAL VELOCITIES



Scheme of aerosol effects on precipitation




Aerosol concentration

Conclusions:

Particulate air pollution acts to delay conversion of cloud water into precipitation. This is manifested in different ways under different circumstances:

- Orographic precipitation is suppressed, with some downwind compensation.
- Convective precipitation processes are delayed to greater heights in the clouds, respectively delay the downdraft and allowing the clouds to invigorate further. This causes:
 - In dry and unstable conditions: Reduced precipitation due to very low precipitation efficiency.
 - In tropical and moist subtropical conditions: Enhanced storm vigor overcompensates for the reduced precipitation efficiency.





1E. Ukiah

2E. Lake Spaulding

3E. Bowman

4E. Boca

5A. San Francisco

6A. Sacramento


7A. Pacific House


8A. Cluster of snow packs in the divide line downwind


to Sacramento 9A.Woodfords

10B. Fresno











Cluster of stations: Judean Hills / Judea plains Ending / Starting ratio = 1.17 / 1.38 = 0.85

